

— ABLYBIO, Help Your Research



# PRKRA Rabbit mAb

货号: **AYM31241**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	34kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,PC-12,Mouse testis
细胞定位	Cytoplasm,perinuclear region
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PRKRA.
------	--

## 靶点信息

研究背景	This gene encodes a protein kinase activated by double-stranded RNA which mediates the effects of interferon in response to viral infection. Mutations in this gene have been associated with dystonia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2008]
基因ID	8575
基因名	PRKRA
Swiss	O75569 ( <a href="https://www.uniprot.org/uniprotkb/O75569/entry">https://www.uniprot.org/uniprotkb/O75569/entry</a> )
别名	PRKRA,PRKRA Rabbit mAb,PKR-associated protein X,PKR-associating protein X,Protein activator of the interferon-induced protein kinase,Protein kinase,interferon-inducible double-stranded RNA-dependent activator,PACT,RAX

## 产品验证



Western blot analysis of PRKRA expressed in 293T,PC-12,Mouse testis using PRKRA Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))