

GCSF Receptor (CD114) Rabbit mAb

货号: **AYM31254**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	85kDa/86kDa/92kDa/95kDa
实测分子量	92kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-937,293T
细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human GCSF Receptor (CD114).
序列	CSPNRKNPLWSPVDPDAHSSLGSWVPTIMEEDAFQLPGLGTPPITKLTVLEEDEKKVPWESHNSSETCGLPTLVQTYVLQ GDPRAVSTQPQSQSGTSDQV

靶点信息

研究背景	The protein encoded by this gene is the receptor for colony stimulating factor 3, a cytokine that controls the production, differentiation, and function of granulocytes. The encoded protein, which is a member of the family of cytokine receptors, may also function in some cell surface adhesion or recognition processes. Alternatively spliced transcript variants have been described. Mutations in this gene are a cause of Kostmann syndrome, also known as severe congenital neutropenia.
基因ID	1441

基因名	CSF3R
Swiss	Q99062
别名	CSF3R;CD114;GCSFR;SCN7

产品验证



Western blot analysis of GCSF Receptor (CD114) expressed in U-937,293T using GCSF Receptor (CD114) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn