

# HIF-1 beta Rabbit mAb

货号: **AYM31341**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	87kDa
实测分子量	87kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human HIF-1 beta .
序列	LAPRQQQQTELDMVPGRDGLASYNHSQVVQPVTTTGPEHSPLEKSDGLFAQDRDPRFSEIYHNINADQSKGISSSTV PATQQLFSQGNTPPTPRPAENFRNSGLAPPVTIVQPSASAGQMLAQISRHSNPTQGATPTWPTTRSGFSAQQVATQAT AKTRTSQFGVGSFQTPSSFSSMSLPGAPTASPGAAAYPSLTNRGSNFAPETGQTAGQFQTRTAEGVGVWPQWQQQP HHRSSSSEQHVQPPAQQPGQPEVFQEMLSMLGDQSNNSYNNEEFPDLTMFPPFSE


## 靶点信息

研究背景	This gene encodes a protein containing a basic helix-loop-helix domain and two characteristic PAS domains along with a PAC domain. The encoded protein binds to ligand-bound aryl hydrocarbon receptor and aids in the movement of this complex to the nucleus, where it promotes the expression of genes involved in xenobiotic metabolism. This protein is also a co-factor for transcriptional regulation by hypoxia-inducible factor 1. Chromosomal translocation of this locus with the ETV6 (ets variant 6) gene on chromosome 12 have been described in leukemias. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2013]
基因ID	405
基因名	ARNT
Swiss	P27540
别名	ARNT;HIF-1-beta;HIF-1beta;HIF1-beta;HIF1B;HIF1BETA;TANGO;bHLHe2

## 产品验证



Immunohistochemical analysis of paraffin-embedded human colon, using HIF-1 beta Antibody .



Western blot analysis of HIF-1 beta expressed in HeLa using HIF-1 beta Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 1 20s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)