

STK39 Rabbit mAb

货号: **AYM31351**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	68kDa
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,SH-SY5Y,Rat testis,Mouse testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human STK39.
序列	SQEKSRRVKEENPEIAVSASTIPEQIQLSLVHDSQGPPNANEDYREASSCAVNLVLRRLNRSRKELNDIRFEFTPGRDADGV SQELFSAGLVDGHDVVIVAANLQKIVDDPKALKTLTFKLAGCDGSEIPDEVKLIQFAQLSVS

靶点信息

研究背景	This gene encodes a serine/threonine kinase that is thought to function in the cellular stress response pathway. The kinase is activated in response to hypotonic stress, leading to phosphorylation of several cation-chloride-coupled cotransporters. The catalytically active kinase specifically activates the p38 MAP kinase pathway, and its interaction with p38 decreases upon cellular stress, suggesting that this kinase may serve as an intermediate in the response to cellular stress. [provided by RefSeq, Jul 2008]
------	---

基因ID	27347
基因名	STK39
Swiss	Q9UEW8
别名	DCHT; PASK; SPAK

产品验证



Western blot analysis of STK39 expressed in Jurkat,SH-SY5Y,Rat testis using STK39 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn