

CD33 Rabbit mAb

货号: **AYM31355**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	25kDa/33kDa/39kDa
实测分子量	65-75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	TF-1, Rat lung
细胞定位	Cell membrane, Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD33.
序列	YYDKNSPVHGYWFRGAIISRDSPVATNKLQVEVQETQGRFRLLGDPSRNNCSLSIVDARRRDNGSYFFRMERGSKYS YKSPQLSVHVTDLTHRPKILIPGTLEPGHSHKNTCSVSWACEQGTPIIFSWLSAAPTSLGPRTHSSVLIITPRPQDHGTNLT CQVKFAGAGVTTERTIQLNVTYVPQNPTTGIFPGDGSQKQETRAGVVH

靶点信息

研究背景	Myeloid cell surface antigen CD33 also known as Sialic acid binding Ig-like lectin 3, CD33 antigen or Siglec-3, is a member of the immunoglobulin superfamily and SIGLEC (sialic acid binding Ig-like lectin) family. This Single-pass type I membrane protein contains 1 Ig-like C2-type (immunoglobulin-like) domain and 1 Ig-like V-type (immunoglobulin-like) domain. CD33/Siglec-3 induces apoptosis in acute myeloid leukemia (in vitro). CD33/Siglec-3 can function as a sialic acid-dependent cell adhesion molecule and that binding can be modulated by endogenous sialoglycoconjugates when CD33 is expressed in a plasma membrane.
------	--

基因ID	945
基因名	CD33
Swiss	P20138
别名	CD33;SIGLEC-3;SIGLEC3;p67

产品验证



Western blot analysis of CD33 expressed in TF-1, Rat lung using CD33 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn