

Septin 8 Rabbit mAb

货号: **AYM31369**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	43kDa/49kDa/51kDa/55kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,HeLa,PC-3(negative),Mouse brain,Mouse heart,Rat brain,Rat lung
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Septin 8.
序列	NKVKETELELKEKERELHEKFEHLKRVHQEKRKVEEKRRELEEEETNAFNRRKAAVEALQSQUALHATSQQPLRDKDKKKN

靶点信息

研究背景	This gene is a member of the septin family of nucleotide binding proteins, originally described in yeast as cell division cycle regulatory proteins. Septins are highly conserved in yeast, Drosophila, and mouse, and appear to regulate cytoskeletal organization. Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid cells. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	23176
基因名	SEPT8
Swiss	Q92599
别名	SEPT8;42980;SEP2;septin-8

产品验证



Western blot analysis of Septin 8 expressed in Mouse brain,Rat brain,K-562 using Septin 8 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn