— ABLYBIO, Help Your Research

# Hsp90 Rabbit mAb

**货号: AYM31393**

## 产品信息

| | |
|---|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC IP FC |
| 推荐浓度 | **WB:** 1:500 - 1:2000<br>**IHC:** 1:50 - 1:200<br>**IF/ICC:** 1:50 - 1:200<br>**IP:** 1:20 - 1:50<br>**FC:** 1:20 - 1:50 |
| 理论分子量 | 90kDa |
| 实测分子量 | 90kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20℃. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,Raji,Mouse liver,Rat liver,Rat brain |
| 细胞定位 | Cell membrane,Cytoplasm,Melanosome |
| 纯化 | Affinity purification |


## 抗原信息

| | |
|---|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human Hsp90 . |


## 靶点信息

| | |
|---|---|
| 研究背景 | The protein encoded by this gene is an inducible molecular chaperone that functions as a homodimer. Th e encoded protein aids in the proper folding of specific target proteins by use of an ATPase activity that is modulated by co-chaperones. Two transcript variants encoding different isoforms have been found for thi s gene. [provided by RefSeq, Jan 2012] |
| 基因ID | 3320,3326 |
| 基因名 | HSP90AA1,HSP90AB1 |
| Swiss | P07900P08238 (https://www.uniprot.org/uniprotkb/P07900P08238/entry) |
| 别名 | Hsp90,Hsp90 Rabbit mAb,HSP90AA1,HSP90AB1 |


## 产品验证





Western blot analysis of Hsp90 expressed in Mouse liver,Rat liver,HeLa using Hsp90 Rabbit mAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.




Immunohistochemical analysis of paraffin-embedded Rat spleen, using the Antibody at 1:300 dilution.




Immunohistochemical analysis of paraffin-embedded Human Hodgkin's lymphoma, using the Antibody at 1:100 dilution.




Immunohistochemical analysis of paraffin-embedded Human glioblastoma, using the Antibody at 1:100 dilution.



Immunohistochemical analysis of paraffin-embedded Human esophageal carcinoma, using the Antibody at 1:100 dilution.



Immunohistochemical analysis of paraffin-embedded Mouse pancreas, using the Antibody at 1:300 dilution.



Immunohistochemical analysis of paraffin-embedded human liver carcinoma, using Hsp90 Antibody .

**实验步骤**

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)