

Gelsolin Rabbit mAb

货号: **AYM31400**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	80kDa/81kDa/85kDa
实测分子量	86kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,SW480
细胞定位	Cytoplasm,Secreted,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Gelsolin.
序列	PMIYKGGTSREGGQTAPASTRLFQVRANSAGATRAVEVLPKAGALNSNDAFVLKTPSAAYLWVGTGASEAEKTGAQELL RVLRAQPVQVAEGSEPDGFWEALGGKAAAYRTSPRLKDKKMDAHPRLFACSNKIGRFVIEEVPGELMQEDLATDDVMLL DTWDQVFVWVGKDSQEEKTEALTSAKRYIETDPANRDRRTPITVVVKQGFEPSPFVGWFLGWDDDDYWSVDPLDRAMA ELAA

靶点信息

研究背景	The protein encoded by this gene binds to the 'plus' ends of actin monomers and filaments to prevent monomer exchange. The encoded calcium-regulated protein functions in both assembly and disassembly of actin filaments. Defects in this gene are a cause of familial amyloidosis Finnish type (FAF). Multiple transcript variants encoding several different isoforms have been found for this gene.
------	--

基因ID	2934
基因名	GSN
Swiss	P06396
别名	GSN;ADF;AGEL;gelsolin

产品验证



Western blot analysis of Gelsolin expressed in U-251MG,SW480 using Gelsolin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn