

— ABLYBIO, Help Your Research



# ADAR1 Rabbit mAb

货号: **AYM31428**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	150kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T
细胞定位	Cytoplasm,Nucleus,Nucleus,nucleolus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ADAR1.
------	--

## 靶点信息

研究背景	This gene encodes the enzyme responsible for RNA editing by site-specific deamination of adenosines. This enzyme destabilizes double-stranded RNA through conversion of adenosine to inosine. Mutations in this gene have been associated with dyschromatosis symmetrica hereditaria. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2010]
基因ID	103
基因名	ADAR
Swiss	P55265
别名	ADAR1,ADAR1 Rabbit mAb,ADAR,136 kDa double-stranded RNA-binding protein,Interferon-inducible protein 4,K88DSRBP,DSRAD,G1P1,IFI4

## 产品验证



Western blot analysis of ADAR1 expressed in HeLa,293T using ADAR1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))