

FGF19 Rabbit mAb

货号: **AYM31440**

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 24kDa |
| 实测分子量 | 36kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HT-29 |
| 细胞定位 | Secreted |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human FGF19. |
| 序列 | LAFSDAGPHVHYGWGDPRLRLHLYTSGPHGLSSCFRLRIRADGVVDCARGQSAHSLLEIKAVLRRTVAIKGVHHSVRYLCMG ADGKMQLLQYSEEDCAFEIEIRPDGYNVYRSEKHRLPVLSLSSAKQRQLYKNRGLPLSHFLPMLPMVPEEPEDLRGHLES DMFSSPLETDSMDPFGLVTGLEAVRSPSFEK |

靶点信息

| | |
|------|--|
| 研究背景 | The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes including embryonic development cell growth, morphogenesis, tissue repair, tumor growth and invasion. This growth factor is a high affinity, heparin dependent ligand for FGFR4. Expression of this gene was detected only in fetal but not adult brain tissue. Synergistic interaction of the chick homolog and Wnt-8c has been shown to be required for initiation of inner ear development. |
|------|--|

| | |
|-------|--------|
| 基因ID | 9965 |
| 基因名 | FGF19 |
| Swiss | O95750 |
| 别名 | FGF19 |

产品验证



Western blot analysis of FGF19 expressed in HT-29 using FGF19 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn