

ZMYND8 Rabbit pAb

货号: **AYP10343**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	66kDa/82kDa/109-138kDa
实测分子量	165kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87 MG,LO2,MCF-7,A375
细胞定位	cytoplasm,Golgi apparatus,nucleoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 959-1188 of human ZMYND8 (NP_898868.1).
序列	LELTMAEMRQSLEQERDRLIAEVKKQLELEKQQAQVDETKKKQWCANCKKEAIFYCCWNTSYCDYPCQQAHWPEHMKSC TSATAPQQEADAENVNTEETLNKSSQSSSSTQSAPSETASASKEKETSAEKSKESGSTLDLSGSRETPSSILLGNSQGSVS KRCQKQPAYAPTTTDDHQHPNYPYPAQKYHSRSNKSSWSSSDEKRGSTRSDHNTSTSTKSLLPKESRLDTFWD

靶点信息

研究背景	The protein encoded by this gene is a receptor for activated C-kinase (RACK) protein. The encoded protein has been shown to bind in vitro to activated protein kinase C beta I. In addition, this protein is a cutaneous T-cell lymphoma-associated antigen. Finally, the protein contains a bromodomain and two zinc fingers, and is thought to be a transcriptional regulator. Multiple transcript variants encoding several different isoforms have been found for this gene.
------	--

基因ID	23613
基因名	ZMYND8
Swiss	Q9ULU4
别名	ZMYND8;PRKCBP1;PRO2893;RACK7

产品验证



Western blot analysis of ZMYND8 expressed in U-87 MG, LO2, MCF-7 using ZMYND8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn