

Flotillin 1 Rabbit pAb

货号: **AYP10393**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:500 - 1:1000
理论分子量	42kDa/47kDa
实测分子量	47KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,A-549,Mouse brain,Rat brain
细胞定位	Cell membrane,Endosome,Melanosome,Membrane,Peripheral membrane protein,caveola
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 155-427 of human Flotillin 1 (NP_005794.1).
序列	HDDQDYLSLHGKARTAQVQKDARIGEAEAKRDAGIREAKAKQEKVSAQYLSEIEMAKAQRDYELKKAAYDIEVNTRRAQADLAYQLQVAKTKQQIEEQRVQVQVVERAQQVAVQEIEIARREKELEARVRKPAAERYKLERLAEAEKSQLIMQAEAEAASVRMRGEAEFAIGARARAEAEQMAKKAEAFQLYQEAQDLMLLEKLPQVAEEISGPLTSANKITLVSSSGTGMGAAKVTGEVLDILTRLPESVERLTGVSISQVNHKPLRTA

靶点信息

研究背景	This gene encodes an protein that localizes to the caveolae, which are small domains on the inner cell membranes. This protein plays a role in vesicle trafficking and cell morphology. Alternative splicing results in multiple transcript variants.
基因ID	10211
基因名	FLOT1
Swiss	O75955
别名	FLOT1

产品验证



Western blot analysis of Flotillin 1 expressed in Mouse brain,Rat brain,293T using Flotillin 1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn