

CCT8 Rabbit pAb

货号: **AYP10400**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rattus norvegicus
应用	WB IHC
推荐浓度	WB: 1:1000 - 1:2000 IHC: 1:50 - 1:200
理论分子量	51kDa/57kDa/59kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	B cells,LO2,293T,HeLa,Mouse brain,Mouse liver,Mouse lung,Rat brain
细胞定位	Cytoplasm,centrosome,cilium basal body,cytoskeleton,microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 248-497 of human CCT8 (NP_001269837.1).
序列	DMALHYANKYNIMLVRLNSKWDLRRLCKTVGATALPRLTPPVLEEMGHCDVSVYLSEVGDQVVFVKHEKEDGAISTIVLRG STDNLMDIERAVDDGVNTFKVLTRDKRLVPGGGATEIELAKQITSYGETCPGLEQYAIKKFAEAFEIPRALAENSGVKAN EVISKLYAVHQEGNKVGLDIEAEVPAVKDMLEAGILDTYLGKYWAIKLATNAAVTVLRVDQIIMAKPAGGPKPPSGKKDW DDDQND

靶点信息

研究背景	This gene encodes the theta subunit of the CCT chaperonin, which is abundant in the eukaryotic cytosol and may be involved in the transport and assembly of newly synthesized proteins. Alternative splicing results in multiple transcript variants of this gene. A pseudogene related to this gene is located on chromosome 1.
------	--

基因ID	10694
基因名	CCT8
Swiss	P50990
别名	CCT8;C21orf112;Cctq;D21S246;PRED71

产品验证



Western blot analysis of CCT8 expressed in Mouse brain,Rat brain,B cells using CCT8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn