

CD45 Rabbit pAb

货号: **AYP10604**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Human chronic myeloid leukemia cells , Mus musculus IF: Human chronic myeloid leukemia cells , Mus musculus , Rattus norvegicus WB: Mus musculus FC: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	130kDa/147kDa
实测分子量	180-250KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,MCF7
细胞定位	Membrane,Membrane raft,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 26-577 of human CD45 (P08575).
序列	QSPTPSPTGLTTAKMPSVPLSSDPLPHTHTTAFSPASTFERENDFSETTSLSPDNTSTQVSPDSLNASAFNTTGVSSVQT PHLPHTADSQTSPSAGTDTQTFSGSAANAKLNPTPGSNAISDVPGERSTASTFPTDPVSPLTTTTLSLAHHSSAALPARTSNT TITANTS DAYLNASETTTLSPSGSAVISTTTIATTPSKPTCDEKYANITVDYLYNKETKLFTAKLNVNENVECGNNTCTNNEVH NLTECKNASVSISHNSCTAPDKTLILDVPPGVEKFLHDCTQVEKADTTICLKWKNIETFTCDTQNITYRFQCGNMIFDNKEI KLENLEPEHEYKCDSEILYNNHKFTNASKIITDFGSPGEPQIIFCRSEAAHQGVITWNPPQRSFHNFTLCYIKETEKDCLNLD KNLIKYDLQNLKPYTKYVLSLHAYIIAKVQRNGSAAMCHFTTKSAPPSQVWNMTVSMTSDNSMHVKRPPRDRNGPHERY HLEVEAGNTLVRNESHKNCDFRVKDLQYSTDYTFKAYFHNGDYGPGEFILHHSTSYNSK

靶点信息

研究背景	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported.
基因ID	5788
基因名	PTPRC
Swiss	P08575
别名	B220;CD45;CD45R;GP180;L-CA;LCA;LY5;T200;PTPRC

产品验证



Western blot analysis of CD45 expressed in Jurkat, MCF7 using CD45 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn