

Acetyl-Histone H3-K18 Rabbit pAb

货号: **AYP10616**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Saccharomyces cerevisiae , Homo sapiens , Monascus ruber CHIP: Homo sapiens
应用	WB IHC IP CHIP
推荐浓度	WB: 1:100 - 1:500 IHC: 1:200 - 1:500 IF/ICC: 1:500 - 1:1000 IP: 1:200 - 1:500 ChIP: 1:50 - 1:100
理论分子量	15kDa
实测分子量	17KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Chromosome,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic acetylated peptide around K18 of human H3 (NP_003520.1).
序列	PRKQL

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.
基因ID	8356
基因名	
Swiss	P68431
别名	HIST1H3J;H3/j;H3FJ

产品验证



Western blot analysis of Acetyl-Histone H3-K18 expressed in HeLa using Acetyl-Histone H3-K18 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn