

SARS2 Rabbit pAb

货号: **AYP10673**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens IF: Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	58kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse liver,Mouse kidney
细胞定位	Mitochondrion matrix
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 289-518 of human SARS 2 (NP_060297.1).
序列	ARFKDLNLAGTAEVGLAGYFMDHTVAFRDLPVRMVCSSSTCYRAETNTGQEPRGLYRVHHFTKVEMFGVTGPGLEQSSQL LEEFSLQMEILTELGLHFRVLDMPQTQELGLPAYRKFDIEAWMPGRGRFGEVTSASNCTDFQSRRLHIMFQTEAGELQFAH TVNATACAVPRLLIALLESNQKDGSVLPPALQSYLGTDRITAPTHVPLQYIGPNQPRKPGPLGQPAVS

靶点信息

研究背景	This gene encodes the mitochondrial seryl-tRNA synthetase precursor, a member of the class II tRNA synthetase family. The mature enzyme catalyzes the ligation of Serine to tRNA(Ser) and participates in the biosynthesis of selenocysteinyl-tRNA(sec) in mitochondria. The enzyme contains an N-terminal tRNA binding domain and a core catalytic domain. It functions in a homodimeric form, which is stabilized by tRNA binding. This gene is regulated by a bidirectional promoter that also controls the expression of mitochondrial ribosomal protein S12. Both genes are within the critical interval for the autosomal dominant deafness locus DFNA4 and might be linked to this disease. Multiple transcript variants encoding different isoforms have been identified for this gene.
基因ID	54938
基因名	SARS2
Swiss	Q9NP81
别名	SARS2;SARS;SARSM;SERS;SYS;SerRS;SerRSmt

产品验证



Western blot analysis of SARS2 expressed in HepG2, Mouse liver, Mouse kidney using SARS2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn