

— ABLYBIO, Help Your Research



KDM3A Rabbit pAb

货号: **AYP10674**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens IF: Mus musculus IHC: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	147kDa
实测分子量	147KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F,HAP1
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 350-450 of human KDM3A (NP_060903.2).
------	--

靶点信息

研究背景	This gene encodes a zinc finger protein that contains a jumonji domain and may play a role in hormone-dependent transcriptional activation. Alternative splicing results in multiple transcript variants.
基因ID	55818
基因名	KDM3A
Swiss	Q9Y4C1 (https://www.uniprot.org/uniprotkb/Q9Y4C1/entry)
别名	KDM3A,JHDM2A,JHMD2A,JMJD1,JMJD1A,TSGA,KDM3A Rabbit pAb,JmjC domain-containing histone demethylation protein 2A,Jumonji domain-containing protein 1A,[histone H3]-dimethyl-L-lysine(9) demethylase 3A, KIAA0742

产品验证



Western blot analysis of KDM3A expressed in 293F,HAP1 using KDM3A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)