

# KDM3A Rabbit pAb

货号: **AYP10674**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus , Homo sapiens <b>IF:</b> Mus musculus <b>IHC:</b> Homo sapiens
应用	<a href="#">WB</a> <a href="#">IHC</a> <a href="#">IF/ICC</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	147kDa
实测分子量	147KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F,HAP1
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 350-450 of human KDM3A (NP_060903.2).
序列	LNTKSEALRTKPDVCKAGLLSKSSQIGTGLKILTEPKGSCTQPKTNTDQENRLESVPQALTGLPKECLPTKASSKAELEIANPPELQKHLEHAPSPSDVS

## 靶点信息

研究背景	This gene encodes a zinc finger protein that contains a jumonji domain and may play a role in hormone-dependent transcriptional activation. Alternative splicing results in multiple transcript variants.
------	---

基因ID	55818
基因名	KDM3A
Swiss	Q9Y4C1
别名	KDM3A;JHDM2A;JHMD2A;JMJD1;JMJD1A;TSGA

## 产品验证



Western blot analysis of KDM3A expressed in 293F, HAP1 using KDM3A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)