

— ABLYBIO, Help Your Research



# AMH Rabbit pAb

货号: **AYP10689**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:100 - 1:500 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	59kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,Mouse testis,Rat liver,Rat testis
细胞定位	Secreted
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 360-560 of human AMH (NP_000470.3).
------	---

## 靶点信息

研究背景	Anti-Mullerian hormone is a member of the transforming growth factor-beta gene family which mediates male sexual differentiation. Anti-Mullerian hormone causes the regression of Mullerian ducts which would otherwise differentiate into the uterus and fallopian tubes. Some mutations in the anti-Mullerian hormone result in persistent Mullerian duct syndrome.
基因ID	268
基因名	AMH
Swiss	P03971 ( <a href="https://www.uniprot.org/uniprotkb/P03971/entry">https://www.uniprot.org/uniprotkb/P03971/entry</a> )
别名	AMH,MIF,MIS,AMH Rabbit pAb,Muellerian-inhibiting factor,Muellerian-inhibiting substance

## 产品验证



Western blot analysis of AMH expressed in Mouse testis,Rat liver,SKOV3 using AMH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)