

# NME3 Rabbit pAb

货号: **AYP10750**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:100 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	19kDa
实测分子量	19kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,Raji,Mouse kidney,Mouse liver
细胞定位	cytosol
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-169 of human NME3 (NP_002504.2).
序列	AHERTFLAVKPDGVQRRLLVGEIVRRFERKGFKLVALKLVQASEELLREHYAELRERPFYGRVLVKYMASGPVVAMVWQGLD VVRTSRALIGATNPADAPPGTIRGDFCIEVGKNIHGSDSVESARREIALWFRADLLCWEDSAGHWLYE

## 靶点信息

研究背景	Major role in the synthesis of nucleoside triphosphates other than ATP. The ATP gamma phosphate is transferred to the NDP beta phosphate via a ping-pong mechanism, using a phosphorylated active-site intermediate. Probably has a role in normal hematopoiesis by inhibition of granulocyte differentiation and induction of apoptosis.
------	---

基因ID	4832
基因名	NME3
Swiss	Q13232
别名	NME3;DR-nm23;NDPK-C;NDPKC;NM23-H3;NM23H3;c371H6.2

## 产品验证



Western blot analysis of NME3 expressed in Mouse kidney,THP-1,Raji using NME3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)