

# CNTRL Rabbit pAb

货号: **AYP10757**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	56kDa/116kDa/205kDa/268kDa
实测分子量	280KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse lung,Mouse spleen
细胞定位	Cytoplasm,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 250-500 of human CNTRL (NP_008949.4).
序列	LEGQPVTTQDRQEAFERFSLEEVEERLERDLEKKMIETEELKSKQTRFLEEIKNQDKLNKSLKEEAMLQKQSCEELKSDLNTK NELLKQKTIELTRACQKQYELEQELAFYKIDAKFEPLNYPSEYAEIDKAPDESPYIGKSRKRNMFATESYIIDSQAQAVQIKK MEPDEQLRNDHMNLRGHTPLDTQLEDKEKKISAAQTRLSELHDEIEKAEQQILRATEEFKQLEEAIQLKKISEAGKDLLYKQL SG

## 靶点信息

研究背景	This gene encodes a centrosomal protein required for the centrosome to function as a microtubule organizing center. The gene product is also associated with centrosome maturation. One version of stem cell myeloproliferative disorder is the result of a reciprocal translocation between chromosomes 8 and 9, with the breakpoint associated with fibroblast growth factor receptor 1 and centrosomal protein 1.
------	--

基因ID	11064
基因名	CNTRL
Swiss	Q7Z7A1
别名	CNTRL;CEP1;CEP110;FAN;bA165P4.1

## 产品验证



Western blot analysis of CNTRL expressed in Raji, Mouse lung, Mouse spleen using CNTRL Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)