

TEX2 Rabbit pAb

货号: **AYP10759**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	125kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse liver
细胞定位	Membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 915-1134 of human TEX 2 (NP_060939.3).
序列	LETKMNLTKLGKEPLVEALKVGEIGKEGCRPRAFCLADSDEESSAGSSEEDDAPEPSGGDKQLLPGAEGYVGGHRTSKI MRFVDKITKSKYFQKATETEFIKKKIEEVSNTPLLLTVEVQECRGLAVNIPPPPTDRVWYGFRKPPHVELKARPKLGEREVTL VHVTDWIEKKLEQEFQKVFVMPNMDVYITIMHSAMDPRSTSCLLKDPPVEAADQP

靶点信息

研究背景	During endoplasmic reticulum (ER stress or when cellular ceramide levels increase, may induce contacts between the ER and medial-Golgi complex to facilitate non-vesicular transport of ceramides from the ER to the Golgi complex where they are converted to complex sphingolipids, preventing toxic ceramide accumulation.
基因ID	55852

基因名	TEX2
Swiss	Q8IWB9
别名	TEX2;HT008;TMEM96

产品验证



Western blot analysis of TEX2 expressed in Mouse kidney, Mouse liver using TEX2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn