

EIF3J Rabbit pAb

货号: **AYP10770**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:200 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	22kDa/23kDa/29kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,HeLa,Jurkat,A-549,U-87MG,LO2,Mouse testis,Mouse liver
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-258 of human EIF3J (N P_003749.2).
序列	MAAAAAAGDSDSWDADAFSVEDPVRKVGGGGTAGGDRWEGEDEDEDVKNWDDDDDEKKEEAEVKPEVKISEKK KIAEKIKEKERQKKRQEEIKRLEEPKVLTPPEQLADKLRLLKQLQEESDLAKETFGVNNNAVYGDAMNPSSRDDFTE FGKLLKDKITQYKSLYYASFLEVLVRDVCISLEIDDLKKITNSLTVLCSEKQKQKQSKAKKKKKGVPPGGGLKATMKDDL ADYGGYDGGYVQDYEDFM

靶点信息

研究背景	This gene encodes a core subunit of the eukaryotic initiation factor 3 complex, which participates in the initiation of translation by aiding in the recruitment of protein and mRNA components to the 40S ribosome. There are pseudogenes for this gene on chromosomes 1, 3, and 9. Alternative splicing results in multiple transcript variants encoding different isoforms.
------	--

基因ID	8669
基因名	EIF3J
Swiss	O75822
别名	EIF3J;EIF3S1;eIF3-alpha;eIF3-p35

产品验证



Western blot analysis of EIF3J expressed in Mouse testis,SKOV3,HeLa using EIF3J Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn