

# DIS3L Rabbit pAb

货号: **AYP10807**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	86kDa/105kDa/111kDa/120kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HeLa,Mouse testis,Rat thymus
细胞定位	Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human DIS3L ( NP_588616.1).
序列	MQTACQAVQHQRGRRQYNKLRNLLKДАРHDCILFANEFQCCYLPRERGESMEKWQTRSIYNAAVWYYHHCQDRMPIV MVTEDEEAIQQYGSETEGVFVITFKNYLDNFWPDLKAAHELCDSILQSRRERENESQESHGKEYPEHLPLEVLEAGIKSGRY IQGILNVNKHRAQIEAFVRLQGASSKSDSLVSDILIHGMKARNRSIHGDVVVVVLLPKNEWKGRTVALCENDCDDKASGE SPSEPMPTGRVVGILQKNWRDYVVTFPSKEEVQSQGKNAQKILVTPWDYRIPKIRISTQQ

## 靶点信息

研究背景	The cytoplasmic RNA exosome complex degrades unstable mRNAs and is involved in the regular turnover of other mRNAs. The protein encoded by this gene contains 3'-5' exoribonuclease activity and is a catalytic component of this complex.
------	--

基因ID	115752
基因名	DIS3L
Swiss	Q8TF46
别名	DIS3L;DIS3L1

## 产品验证



Western blot analysis of DIS3L expressed in Mouse testis,Rat thymus,U-87MG using DIS3L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)