

TRAPPC6A Rabbit pAb

货号: **AYP10829**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	11kDa/12kDa/17kDa/18kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Rat kidney,Rat liver
细胞定位	Endoplasmic reticulum,Golgi apparatus,cis-Golgi network
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-173 of human TRAPPC 6A (NP_077013.1).
序列	MADTVLFEFLHTEMVAELWAHDPDPGPGVSAGLRGEEAGATKGQKMSLSVLEGMGFRVQGALGERLPRETLAFREELD VLKFLCKDLWVAVFQKQMDSLRNTNHQGTIVLQDNSFPLLLPMASGLQYLEEAPKFLAFTCGLLRGALYTLGIESVVTASVA ALPVCKFQVVIPKS

靶点信息

研究背景	This gene encodes a component of the trafficking protein particle complex, which tethers transport vesicles to the cis-Golgi membrane. Loss of expression of the related gene in mouse affects coat and eye pigmentation, suggesting that the encoded protein may be involved in melanosome biogenesis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
------	--

基因ID	79090
基因名	TRAPPC6A
Swiss	O75865
别名	TRAPPC6A;TRS33

产品验证



Western blot analysis of TRAPPC6A expressed in Mouse kidney,Rat kidney,Rat liver using TRAPPC6A R abbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn