

— ABLYBIO, Help Your Research



PGC1 α Rabbit pAb

货号: **AYP11008**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: mouse cells , Rattus norvegicus WB: Rabbit cerebral microvascular endothelial cells , Homo sapiens , Mus musculus , Rattus norvegicus , Gallus gallus , Capra hircus , Hordeum vulgare IHC: Mus musculus , Homo sapiens , Rattus norvegicus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	14kDa/30kDa/31kDa/33kDa/77kDa/89kDa/91kDa
实测分子量	91KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,Mouse heart,Mouse skeletal muscle,Rat heart
细胞定位	Cytoplasm,Nucleus,PML body
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 610-798 of human PGC1 α (NP_037393.1).
------	---

靶点信息

研究背景	The protein encoded by this gene is a transcriptional coactivator that regulates the genes involved in energy metabolism. This protein interacts with PPARgamma, which permits the interaction of this protein with multiple transcription factors. This protein can interact with, and regulate the activities of, cAMP response element binding protein (CREB) and nuclear respiratory factors (NRFs). It provides a direct link between external physiological stimuli and the regulation of mitochondrial biogenesis, and is a major factor that regulates muscle fiber type determination. This protein may be also involved in controlling blood pressure, regulating cellular cholesterol homeostasis, and the development of obesity.
基因ID	10891
基因名	PPARGC1A
Swiss	Q9UBK2 (https://www.uniprot.org/uniprotkb/Q9UBK2/entry)
别名	PPARGC1A,LEM6,PGC-1(alpha),PGC-1alpha,PGC-1v,PGC1,PGC1A,PPARGC1,PPARG coactivator 1 alpha,PGC 1 alpha,PGC1 α Rabbit pAb,Ligand effect modulator 6

产品验证



Western blot analysis of PGC1 α expressed in Mouse heart,Rat heart,HeLa using PGC1 α Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)