

RIPK3 Rabbit pAb

货号: **AYP11029**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: human kidney cancer cells , Mus musculus , Ctenopharyngodon idellus , Gallus gallus , Homo sapiens , Rattus norvegicus , Sus scrofa , Marine Streptomyces sp IF: Mus musculus ICH: Mus musculus ELISA: Mus musculus IHC: Gallus gallus , Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	25kDa/27kDa/56kDa
实测分子量	46-62KDa (M)
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa(Negative control),L929
细胞定位	Cell membrane,Cytoplasm,Mitochondrion,cytosol
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human RIP3 (NP_006862.2).
序列	GSLSGLLQSQCPRPWPLLCRLLKEVVLGMFYLDQNPVLLHRDLKPSNVLLDPELHVKLADFGGLSTFQGGSQSGTGSGEPGGTLGYLAPELFFVNVNRKAST

靶点信息

研究背景	The product of this gene is a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The encoded protein is predominantly localized to the cytoplasm, and can undergo nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-I signaling complex, and can induce apoptosis and weakly activate the NF-kappaB transcription factor.
基因ID	11035
基因名	RIPK3
Swiss	Q9Y572
别名	RIPK3;RIP3

产品验证



Western blot analysis of RIPK3 expressed in HeLa(Negative control),L929 using RIPK3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn