

# Phospho-I $\kappa$ B $\alpha$ -S32/36 Rabbit pAb

货号: **AYP11044**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens,Mus musculus , Mus musculus , Homo sapiens , Melodinus cochinchinensis , Rattus norvegicus , Schisandra chinensis , Gallus gallus , Cavia porcellus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:100 - 1:500
理论分子量	35kDa
实测分子量	40KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic phosphorylated peptide around S32 & S36 of human I $\kappa$ B $\alpha$ (NP_065390.1).
序列	SGLDS

## 靶点信息

研究背景	This gene encodes a member of the NF-kappa-B inhibitor family, which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal dominant disease.
基因ID	4792

基因名	NFKBIA
Swiss	P25963
别名	IKBA;MAD-3;NFKBI;IKB alpha;NFKBIA;EDAID2

## 产品验证



Western blot analysis of Phospho-IkB $\alpha$ -S32/36 expressed in HeLa using Phospho-IkB $\alpha$ -S32/36 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)