

— ABLYBIO, Help Your Research



Fas Rabbit pAb

货号: **AYP11118**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Rattus norvegicus , Homo sapiens , Anatinae ELISA: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	9kDa/11kDa/14kDa/16kDa/24kDa/35kDa/37kDa
实测分子量	40-50KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HT-1080,HepG2,HeLa
细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 61-335 of human Fas (N P_000034.1).
------	---

靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains a death domain. It has been shown to play a central role in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. The interaction of this receptor with its ligand allows the formation of a death-inducing signaling complex that includes Fas-associated death domain protein (FADD), caspase 8, and caspase 10. The auto proteolytic processing of the caspases in the complex triggers a downstream caspase cascade, and leads to apoptosis. This receptor has been also shown to activate NF-kappaB, MAPK3/ERK1, and MAPK8/JNK, and is found to be involved in transducing the proliferating signals in normal diploid fibroblast and T cells. Several alternatively spliced transcript variants have been described, some of which are candidates for nonsense-mediated mRNA decay (NMD). The isoforms lacking the transmembrane domain may negatively regulate the apoptosis mediated by the full length isoform.
基因ID	355
基因名	FAS
Swiss	P25445 (https://www.uniprot.org/uniprotkb/P25445/entry)
别名	ALPS1A,APO-1,APT1,CD95,FAS1,FASTM,TNFRSF6,FAS,Fas Rabbit pAb,Apo-1 antigen,Apoptosis-mediating surface antigen FAS,FASLG receptor

产品验证



Western blot analysis of Fas expressed in Jurkat, HT-1080, HepG2 using Fas Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)