

PPAR α Rabbit pAb

货号: **AYP11237**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens , Macaca mulatta RT-PCR: Mus musculus
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	18kDa/52kDa
实测分子量	52KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat heart
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human PPAR α (NP_001001928.1).
序列	MVDTESPLCPLSPLEAGDLESPLSEEFLLQEMGNIQEISQSIGEDSSGSFGFTEYQYLGSCPGSDGVSITDTLSPASSPSSVT YPVVPGSVDESPPGALNIECRICGDKASGYHYGVHACEGCKGFFRRTIRLKLVDKCDRSCKIQKKNRNKQCQYCRFHKCLS VGMSHNAIRFGRMPRSEKAKLKAIEILTCEHDIEDSETADLKSLAKRIYEAYLKNFNMNKVKARVILS

靶点信息

研究背景	Peroxisome proliferators include hypolipidemic drugs, herbicides, leukotriene antagonists, and plasticizers; this term arises because they induce an increase in the size and number of peroxisomes. Peroxisomes are subcellular organelles found in plants and animals that contain enzymes for respiration and for cholesterol and lipid metabolism. The action of peroxisome proliferators is thought to be mediated via specific receptors, called PPARs, which belong to the steroid hormone receptor superfamily. PPARs affect the expression of target genes involved in cell proliferation, cell differentiation and in immune and inflammatory responses. Three closely related subtypes (alpha, beta/delta, and gamma) have been identified. This gene encodes the subtype PPAR-alpha, which is a nuclear transcription factor. Multiple alternatively spliced transcript variants have been described for this gene, although the full-length nature of only two has been determined.
基因ID	5465
基因名	PPARA
Swiss	Q07869
别名	PPAR;NR1C1;hPPAR;PPARalpha;PPAR alpha;PPARA

产品验证



Western blot analysis of PPAR α expressed in Rat heart using PPAR α Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn