

# TLR4 Rabbit pAb

货号: **AYP11241**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Rattus norvegicus , Mus musculus , Homo sapiens <b>IHC:</b> Rattus norvegicus <b>IF:</b> Mus musculus
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	73kDa/91kDa/95kDa
实测分子量	110KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse skeletal muscle,Rat heart,HeLa
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 27-228 of human TLR4 ( NP_612564.1).
序列	EPCVEVVPNITYQCMELNFYKIPDNLPFSTKNLDLSFNPLRHLGYSYFFSFPELQVLDLSRCEIQTIEDGAYQSLSHLSTLILT GNPIQSLALGAFSGLSSLQKLVAVETNLALENFPIGHLKTLKELNVAHNLIQSFKLPEYFSNLTNLEHLDLSSNKIQSIYCTD LRVLHQMPLLNLSLDLSLNPMNFIQPGAFKEIRL

## 靶点信息

研究背景	The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from <i>Drosophila</i> to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor has been implicated in signal transduction events induced by lipopolysaccharide (LPS) found in most gram-negative bacteria. Mutations in this gene have been associated with differences in LPS responsiveness. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	7099
基因名	TLR4
Swiss	O00206
别名	TLR4;ARMD10;CD284;TLR-4;TOLL

## 产品验证



Western blot analysis of TLR4 expressed in Mouse skeletal muscle, Rat heart, HeLa using TLR4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)