

TEAD1 Rabbit pAb

货号: **AYP11317**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Mus musculus , Xenopus laevis Co-IP: Homo sapiens IF: Mus musculus ChIP: Mus musculus IHC: Homo sapiens
应用	WB IHC IF/ICC ChIP
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 ChIP: 1:50 - 1:200
理论分子量	40kDa/47kDa
实测分子量	50KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart,Mouse lung
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 135-215 of human TEAD 1 (NP_068780.2).
序列	AIHNKLGGLPGIPRPTFPGAPGFWPGMIQTGQPGSSQDVKPFVQQAYPIQPAVTAPIPGFEPASAPAPSVPAWQGRSIGTTK

靶点信息

研究背景	This gene encodes a ubiquitous transcriptional enhancer factor that is a member of the TEA/ATTS domain family. This protein directs the transactivation of a wide variety of genes and, in placental cells, also acts as a transcriptional repressor. Mutations in this gene cause Sveinsson's chorioretinal atrophy. Additional transcript variants have been described but their full-length natures have not been experimentally verified.
基因ID	7003
基因名	TEAD1
Swiss	P28347
别名	TEAD1;AA;NTEF-1;REF1;TCF-13;TCF13;TEAD-1;TEF-1

产品验证



Western blot analysis of TEAD1 expressed in Mouse heart, Mouse lung using TEAD1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn