

WNT3A Rabbit pAb

货号: **AYP11394**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Mus musculus WB: Mus musculus , Homo sapiens , Rattus norvegicus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	39kDa/42kDa
实测分子量	39KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U2OS
细胞定位	Secreted,extracellular matrix,extracellular space
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human WNT3A (NP_149122.1).
序列	PVLDKATRESAFVHAIASAGVAVAFVTRSCAEGTAAICGSSRHQSPGKGWKWGGCSEDIIEFGGMVSREFADARENRP DARSAMNRHNNEAGRQAIASHMH

靶点信息

研究背景	The WNT gene family consists of structurally related genes which encode secreted signaling proteins. The se proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It encodes a protein which shows 96% amino acid identity to mouse Wnt3A protein, and 84% to human WNT3 protein, another WNT gene product. This gene is clustered with WNT14 gene, another family member, in chromosome 1q42 region.
------	---

基因ID	89780
基因名	WNT3A
Swiss	P56704
别名	WNT3A

产品验证



Western blot analysis of WNT3A expressed in U2OS using WNT3A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn