

# Phospho-RIP3-T231/S232 Rabbit pAb

货号: AYP11414

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens , Mus musculus <b>ICH:</b> Mus musculus <b>IF:</b> Mus musculus <b>ELISA:</b> Mus musculus
应用	WB
推荐浓度	<b>WB:</b> 1:100 - 1:500
理论分子量	53kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	L929
细胞定位	cytoplasm,cytosol,nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic phosphorylated peptide around T231 & S232 of mouse Ripk3 (NP_064339.2).
序列	DKTSLI

## 靶点信息

研究背景	The product of this gene is a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The encoded protein is predominantly localized to the cytoplasm, and can undergo nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-1 signaling complex, and can induce apoptosis and weakly activate the NF-kappaB transcription factor.
------	--

基因ID	56532
基因名	
Swiss	Q9QZL0
别名	R; Rip3; AW107945; 2610528K09Rik

## 产品验证



Western blot analysis of Phospho-RIP3-T231/S232 expressed in L929 using Phospho-RIP3-T231/S232 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)