

CD163 Rabbit pAb

货号: **AYP11440**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Rattus norvegicus , Homo sapiens WB: Mus musculus IF: Homo sapiens , Mus musculus , Gallus gallus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	121kDa/124kDa/125kDa
实测分子量	145KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart,Mouse thymus,Rat liver
细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 900-1000 of human CD163 (NP_004235.4).
序列	DTLWQCPSSPWEKRLASPSEETWITCDNKIRLQEGPTSCSGRVEIWHGGSWGTVCDSDWLDDAQVVCQQLGCGPALKAFKEAEFGQGTGPIWLNEVKCKG

靶点信息

研究背景	The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages, and may thereby protect tissues from free hemoglobin-mediated oxidative damage. This protein may also function as an innate immune sensor for bacteria and inducer of local inflammation. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.
基因ID	9332
基因名	CD163
Swiss	Q86VB7
别名	CD163;M130;MM130;SCAR11

产品验证



Western blot analysis of CD163 expressed in Mouse heart, Mouse thymus, Rat liver using CD163 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s

实验步骤

访问官网浏览详情: www.ablybio.cn