

— ABLYBIO, Help Your Research



# UHRF1 Rabbit pAb

货号: **AYP11518**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> A172 , Homo sapiens <b>IP:</b> Homo sapiens <b>IHC:</b> Mus musculus
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:100 - 1:500 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:20 - 1:50
理论分子量	89kDa/91kDa
实测分子量	95KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human UHRF1 (NP_001041666.1).
------	--

## 靶点信息

研究背景	This gene encodes a member of a subfamily of RING-finger type E3 ubiquitin ligases. The protein binds to specific DNA sequences, and recruits a histone deacetylase to regulate gene expression. Its expression peaks at late G1 phase and continues during G2 and M phases of the cell cycle. It plays a major role in the G1/S transition by regulating topoisomerase IIalpha and retinoblastoma gene expression, and functions in the p53-dependent DNA damage checkpoint. It is regarded as a hub protein for the integration of epigenetic information. This gene is up-regulated in various cancers, and it is therefore considered to be a therapeutic target. Multiple transcript variants encoding different isoforms have been found for this gene. A related pseudogene exists on chromosome 12.
基因ID	29128
基因名	UHRF1
Swiss	Q96T88 ( <a href="https://www.uniprot.org/uniprotkb/Q96T88/entry">https://www.uniprot.org/uniprotkb/Q96T88/entry</a> )
别名	UHRF1,ICBP90,Np95,RNF106,TDRD22,hNP95,hUHRF1,huNp95,UHRF1 Rabbit pAb,Inverted CCAAT box-binding protein of 90 kDa,Nuclear zinc finger protein Np95,RING finger protein 106,Transcription factor ICBP90,Ubiquitin-like PHD and RING finger domain-containing protein 1

## 产品验证



Western blot analysis of UHRF1 expressed in Mouse brain using UHRF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)