

— ABLYBIO, Help Your Research



SIRT6 Rabbit pAb

货号: **AYP11539**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Mus musculus , Homo sapiens |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 36kDa/39kDa |
| 实测分子量 | 39-45kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,293T,Mouse spleen,Mouse stomach,A-549,LO2,BT-474,HL-60,Mouse heart |
| 细胞定位 | Nucleus,nucleoplasm |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of mouse SIRT 6 (NP_001156902.1). |
|------|--|

靶点信息

| | |
|-------|---|
| 研究背景 | This gene encodes a member of the sirtuin family of NAD-dependent enzymes that are implicated in cellular stress resistance, genomic stability, aging and energy homeostasis. The encoded protein is localized to the nucleus, exhibits ADP-ribosyl transferase and histone deacetylase activities, and plays a role in DNA repair, maintenance of telomeric chromatin, inflammation, lipid and glucose metabolism. Alternative splicing results in multiple transcript variants encoding different isoforms. |
| 基因ID | 50721 |
| 基因名 | SIRT6 |
| Swiss | Q8N6T7 |
| 别名 | SIRT6,SIR2L6,sirtuin 6,SIRT6 Rabbit pAb,NAD-dependent protein deacetylase sirtuin-6,Protein mono-ADP-ribose transferase sirtuin-6,Regulatory protein SIR2 homolog 6,SIR2-like protein 6 |

产品验证



Western blot analysis of SIRT6 expressed in Mouse spleen, HeLa, 293T using SIRT6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)