

ASCL4 Rabbit pAb

货号: **AYP11577**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Rattus norvegicus IF: Rattus norvegicus IHC: Rattus norvegicus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	19kDa
实测分子量	19kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse kidney,Rat liver
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-172 of human ASCL4 (NP_982260.2).
序列	MMETRKAERLALPYSLRTAPLGVPGLPGLPRRDPLRVALRLDAACWEWARSGCARGWQYLPVPLDSAFEPFLRKRNERERQVRVCVNEGVARLRDHLPRELADKRLSKVETLRAAIDYIKHLQELLERQAWGLEGAAGAVPQRRRAECNSDGESKASSAPSPSEPEEGG

靶点信息

研究背景	Basic helix-loop-helix transcription factors, such as ASCL4, are essential for the determination of cell fate and the development and differentiation of numerous tissues (Jonsson et al., 2004 [PubMed 15475265]).[supplied by OMIM, Mar 2008]
------	--

基因ID	121549
基因名	ASCL4
Swiss	Q6XD76
别名	ASCL4;HASH4;bHLHa44

产品验证



Western blot analysis of ASCL4 expressed in Mouse brain, Mouse kidney, Rat liver using ASCL4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn