

— ABLYBIO, Help Your Research



AKR1C1 Rabbit pAb

货号: **AYP11587**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Sus scrofa , Mus musculus IHC: Homo sapiens
应用	WB
推荐浓度	WB: 1:1000 - 1:3000
理论分子量	36kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,22Rv1,U-87MG,Mouse liver
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-323 of human AKR1C1 (NP_001344.2).
------	--

靶点信息

研究背景	This gene encodes a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reaction of progesterone to the inactive form 20-alpha-hydroxy-progesterone. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14.
基因ID	1645
基因名	AKR1C1
Swiss	Q04828 (https://www.uniprot.org/uniprotkb/Q04828/entry)
别名	AKR1C1,2-ALPHA-HSD,20-ALPHA-HSD,C9,DD1,DD1/DD2,DDH,DDH1,H-37,HAKRC,HBAB,MBAB,AKR1C1 Rabbit pAb,20alpha-hydroxysteroid dehydrogenase,3beta-hydroxysteroid 3-dehydrogenase,Chlordecone reductase homolog HAKRC,Dihydrodiol dehydrogenase 1,High-affinity hepatic bile acid-binding protein

产品验证



Western blot analysis of AKR1C1 expressed in Mouse liver, HeLa, 22Rv1 using AKR1C1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)