

Rabbit anti His-Tag pAb

货号: **AYP11618**

产品信息

反应	All
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: Homo sapiens , Danio rerio WB: Pseudosciaena crocea , Homo sapiens , Danio rerio
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200 IP: 1:50 - 1:200
理论分子量	
实测分子量	35KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to His tag.
序列	HHHHHH

靶点信息

研究背景	Consecutive histidine residues (usually 6 to 10 in length) are often inserted into the amino acid sequence of recombinant proteins. The resulting His-tagged proteins can be detected by using His antibodies.
基因ID	
基因名	

Swiss	
别名	His;His tag;His-tag

产品验证



Western blot analysis of His-Tag expressed in 293T using Rabbit anti His-Tag pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn