

— ABLYBIO, Help Your Research



# DDX3X Rabbit pAb

货号: **AYP11644**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> 293T , Mus musculus , Chlorocebus aethiops , Homo sapiens <b>IF:</b> Mus musculus
应用	WB IHC IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:500 - 1:1000
理论分子量	71kDa/73kDa
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Rat liver
细胞定位	Cytoplasm,Mitochondrion outer membrane,Nucleus speckle
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human DDX3X (NP_001347.3).
------	---

## 靶点信息

研究背景	The protein encoded by this gene is a member of the large DEAD-box protein family, that is defined by the presence of the conserved Asp-Glu-Ala-Asp (DEAD) motif, and has ATP-dependent RNA helicase activity. This protein has been reported to display a high level of RNA-independent ATPase activity, and unlike most DEAD-box helicases, the ATPase activity is thought to be stimulated by both RNA and DNA. This protein has multiple conserved domains and is thought to play roles in both the nucleus and cytoplasm. Nuclear roles include transcriptional regulation, mRNP assembly, pre-mRNA splicing, and mRNA export. In the cytoplasm, this protein is thought to be involved in translation, cellular signaling, and viral replication. Misregulation of this gene has been implicated in tumorigenesis. This gene has a paralog located in the nonrecombining region of the Y chromosome. Pseudogenes sharing similarity to both this gene and the DDX3Y paralog are found on chromosome 4 and the X chromosome. Alternative splicing results in multiple transcript variants.
基因ID	1654
基因名	DDX3X
Swiss	O00571 ( <a href="https://www.uniprot.org/uniprotkb/O00571/entry">https://www.uniprot.org/uniprotkb/O00571/entry</a> )
别名	DDX3X,CAP-Rf,DBX,DDX14,DDX3,HLP2,MRX102,DDX3X Rabbit pAb,DEAD box protein 3,X-chromosomal,DEAD box,X isoform,Helicase-like protein 2

## 产品验证



Western blot analysis of DDX3X expressed in Mouse brain,Rat liver using DDX3X Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)