

— ABLYBIO, Help Your Research



Phospho-eIF4EBP1-S65 Rabbit pAb

货号: **AYP11672**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB IP
推荐浓度	WB: 1:500 - 1:1000 IP: 1:50 - 1:100
理论分子量	12kDa
实测分子量	20KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	cytoplasm,cytosol
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around S65 of human Phospho-eIF4EBP1-S65 (NP_004086.1).
------	--

靶点信息

研究背景	This gene encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation.
基因ID	1978
基因名	EIF4EBP1
Swiss	Q13541 (https://www.uniprot.org/uniprotkb/Q13541/entry)
别名	4E-BP1,4EBP1,BP-1,PHAS-I,EIF4EBP1,Phospho-eIF4EBP1-S65 Rabbit pAb,Phosphorylated heat- and acid-stable protein regulated by insulin 1

产品验证



Western blot analysis of Phospho-eIF4EBP1-S65 expressed in 293T using Phospho-eIF4EBP1-S65 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)