

MFF Rabbit pAb

货号: **AYP11696**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rattus norvegicus , Homo sapiens , carp IF: carp IHC: Actinopterygii
应用	WB IHC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200
理论分子量	25kDa/27kDa/32kDa/38kDa
实测分子量	25-35KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat heart
细胞定位	Cytoplasmic vesicle,Mitochondrion outer membrane,Peroxisome,Single-pass type IV membrane protein,secretory vesicle,synaptic vesicle
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human MFF (NP_064579.3).
序列	MSKGTSSDTSGLGRVSRAAFPSPTAAEMAEISRIQYEMEYTEGISQRMRVPEKLVAPPNADLEQGFQEGVNPASVIMQVPERIVVAGNNEDVSFSRPADL

靶点信息

研究背景	This is a nuclear gene encoding a protein that functions in mitochondrial and peroxisomal fission. The encoded protein recruits dynamin-1-like protein (DNM1L) to mitochondria. There are multiple pseudogenes for this gene on chromosomes 1, 5, and X. Alternative splicing results in multiple transcript variants.
------	--

基因ID	56947
基因名	MFF
Swiss	Q9GZY8
别名	MFF;C2orf33;EMPF2;GL004

产品验证



Western blot analysis of MFF expressed in Rat heart using MFF Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn