

— ABLYBIO, Help Your Research



# NDUFA13/GRIM19 Rabbit pAb

货号: **AYP11825**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> C2C12 , Homo sapiens
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	16kDa/24kDa
实测分子量	17KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,PC-3
细胞定位	Matrix side,Mitochondrion inner membrane,Nucleus,Single-pass membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-144 of human NDUFA 13/GRIM19 (NP_057049.5).
------	-----------------------------------------------------------------------------------------------------------------------------

## 靶点信息

研究背景	This gene encodes a subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), which functions in the transfer of electrons from NADH to the respiratory chain. The protein is required for complex I assembly and electron transfer activity. The protein binds the signal transducers and activators of transcription 3 (STAT3) transcription factor, and can function as a tumor suppressor. The human protein purified from mitochondria migrates at approximately 16 kDa. Transcripts originating from an upstream promoter and capable of expressing a protein with a longer N-terminus have been found, but their biological validity has not been determined.
基因ID	51079
基因名	NDUFA13
Swiss	Q9P0J0 ( <a href="https://www.uniprot.org/uniprotkb/Q9P0J0/entry">https://www.uniprot.org/uniprotkb/Q9P0J0/entry</a> )
别名	NDUFA13,B16.6,CDA016,CGI-39,GRIM-19,GRIM19,NDUFA13/GRIM19 Rabbit pAb,Cell death regulatory protein GRIM-19,Complex I-B16.6,Gene associated with retinoic and interferon-induced mortality 19 protein,NADH-ubiquinone oxidoreductase B16.6 subunit

## 产品验证



Western blot analysis of NDUFA13/GRIM19 expressed in 293T, HepG2, PC-3 using NDUFA13/GRIM19 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)