

— ABLYBIO, Help Your Research



KDM3A Rabbit pAb

货号: **AYP11828**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: HeLa,MCF7 , Mus musculus , Homo sapiens , Lithobates sylvaticus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	147kDa
实测分子量	147KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human KDM3A (NP_060903.2).
------	---

靶点信息

研究背景	This gene encodes a zinc finger protein that contains a jumonji domain and may play a role in hormone-dependent transcriptional activation. Alternative splicing results in multiple transcript variants.
------	---

基因ID	55818
基因名	KDM3A
Swiss	Q9Y4C1 (https://www.uniprot.org/uniprotkb/Q9Y4C1/entry)
别名	KDM3A,JHDM2A,JHMD2A,JMJD1,JMJD1A,TSGA,KDM3A Rabbit pAb,JmjC domain-containing histone demethylation protein 2A,Jumonji domain-containing protein 1A,[histone H3]-dimethyl-L-lysine(9) demethylase 3A, KIAA0742

产品验证



Western blot analysis of KDM3A expressed in 293T, Mouse testis using KDM3A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)