

RIPK4 Rabbit pAb

货号: **AYP11883**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	86kDa/91kDa
实测分子量	92KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A375
细胞定位	Cytoplasm,Membrane,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 240-520 of human RIPK 4 (NP_065690.2).
序列	HRPELPPVCRARPRACSHLIRLMQRCWQGDPRVRPTFQEITSETEDLCEKPDDEVKETAHDLVDKSPPEPRSEVVPARLKR ASAPTFDNDYLSSELLSQLDSGVSQAVEGPEELSRSSSESKLPSSGSGKRLSGVSSVDSAFSSRGSLSLSFEREPSTSDLG TTDVQKKKLVDAIVSGDTSKLMKILQPQVDLALDSGASLLHLAVEAGQECAKWLLLNNANPNLSNRRGSTPLHMAVER RVRGVVELLLARKISVNAKDEDQWTALHFAAQNGDESST

靶点信息

研究背景	The protein encoded by this gene is a serine/threonine protein kinase that interacts with protein kinase C-delta. The encoded protein can also activate NFkappaB and is required for keratinocyte differentiation. Th is kinase undergoes autophosphorylation.
------	--

基因ID	54101
基因名	
Swiss	P57078
别名	RIPK4;ANKK2;ANKRD3;DIK;NKRD3;PKK;PPS2;RIP4

产品验证



Western blot analysis of RIPK4 expressed in A375 using RIPK4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn