

— ABLYBIO, Help Your Research



COL4A2 Rabbit pAb

货号: **AYP11892**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus IHC: Rattus norvegicus IF: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	167kDa
实测分子量	200kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NCI-H460,HeLa,Mouse lung,Mouse ovary,Rat lung
细胞定位	Secreted,basement membrane,extracellular matrix,extracellular space
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1486-1712 of human C OL4A2 (NP_001837.2).
------	---

靶点信息

研究背景	This gene encodes one of the six subunits of type IV collagen, the major structural component of basement membranes. The C-terminal portion of the protein, known as canstatin, is an inhibitor of angiogenesis and tumor growth. Like the other members of the type IV collagen gene family, this gene is organized in a head-to-head conformation with another type IV collagen gene so that each gene pair shares a common promoter.
基因ID	1284
基因名	COL4A2
Swiss	P08572 (https://www.uniprot.org/uniprotkb/P08572/entry)
别名	COL4A2,ICH,POREN2,COL4A2 Rabbit pAb

产品验证



Western blot analysis of COL4A2 expressed in Mouse lung,Rat lung,NCI-H460 using COL4A2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)