

— ABLYBIO, Help Your Research



SLC23A2 Rabbit pAb

货号: **AYP11910**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus IF: Mus musculus
应用	WB
推荐浓度	WB: 1:1000 - 1:5000
理论分子量	58kDa/70kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F,Neuro-2a
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-98 of human SLC23A2 (NP_005107.4).
------	--

靶点信息

研究背景	The absorption of vitamin C into the body and its distribution to organs requires two sodium-dependent vitamin C transporters. This gene encodes one of the two required transporters and the encoded protein accounts for tissue-specific uptake of vitamin C. Previously, this gene had an official symbol of SLC23A1.
基因ID	9962
基因名	SLC23A2
Swiss	Q9UGH3 (https://www.uniprot.org/uniprotkb/Q9UGH3/entry)
别名	SLC23A2,NBTL1,SLC23A1,SVCT2,YSPL2,hSVCT2,SLC23A2 Rabbit pAb,Na(+)/L-ascorbic acid transporter 2, Nucleobase transporter-like 1 protein,Sodium-dependent vitamin C transporter 2,Yolk sac permease-like molecule 2,KIAA0238

产品验证



Western blot analysis of SLC23A2 expressed in 293F, Neuro-2a using SLC23A2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)