

PENK Rabbit pAb

货号: AYP11920

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rattus norvegicus , Homo sapiens , Mus musculus IHC: Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:200
理论分子量	30kDa
实测分子量	36KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 25-267 of human PENK (NP_001129162.1).
序列	ECSQDCATCSYRLVRPADINFLACVMECEGKLPKSLKIWETCKELLQLSKPELPQDGTSTLRENSKPEESHLLAKRYGGFMKR YGGFMKKMDELYPMEPEEEANGSEILAKRYGGFMKKDAEEDDSLANSDDLKELLETDGDNRRERSHHQDGSNDNEEVSKR YGGFMRGLKRSPQLEDAKELQKRYGGFMRRVGRPEWMDYQKRYGGFLKRFAEALPSDEEGESYSKEVPEMEKRYG GFMRF

靶点信息

研究背景	This gene encodes a preproprotein that is proteolytically processed to generate multiple protein products . These products include the pentapeptide opioids Met-enkephalin and Leu-enkephalin, which are stored in synaptic vesicles, then released into the synapse where they bind to mu- and delta-opioid receptors to modulate the perception of pain. Other non-opioid cleavage products may function in distinct biological activities.
基因ID	5179
基因名	PENK
Swiss	P01210
别名	PENK;PE;PENK-A

产品验证



Western blot analysis of PENK expressed in THP-1 using PENK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn