

ING4 Rabbit pAb

货号: **AYP11926**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	20kDa/25kDa/28kDa
实测分子量	29kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,Raji,A-549,Mouse spleen
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-249 of human ING4 (N P_001121054.1).
序列	MAAGMYLEHYLDSIENLPFELQRNFMRDLDQRTEDLKAIEDKLATEYMSSARSLSSEEKALLKQIQEAYGKCKEFGDD KVQLAMQTYEMVDKHIRRLDLDLARFEADLKEKQIESDYDSSSSKGGKKGRGRTQKEKKAARARSKGKNSDEEAPKTAQK KLKLVRTSPEYGMPSVTFGSVHPSDVLDMVPDPNEPTYCLCHQVSYGEMIGCDNPDCSIEWFHACVGLTTKPRGKWFC PRCSQERKKK

靶点信息

研究背景	This gene encodes a tumor suppressor protein that contains a PHD-finger, which is a common motif in proteins involved in chromatin remodeling. This protein can bind TP53 and EP300/p300, a component of the histone acetyl transferase complex, suggesting its involvement in the TP53-dependent regulatory pathway. Multiple alternatively spliced transcript variants have been observed that encode distinct proteins.
------	--

基因ID	51147
基因名	ING4
Swiss	Q9UNL4
别名	ING4;my036;p29ING4

产品验证



Western blot analysis of ING4 expressed in Mouse spleen, HeLa, 293T using ING4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn